

GoMRI-sponsored Special Section Articles

summer of 2010 (Graham et al. 2010), and mesocosm trials indicate that phytoplankton taxa such as diatoms, coccidiales and chlorophyll *a* have increased in relative (if not total) abundance in oil-contaminated coastal waters (Gilde and Pincus 2012). Changes in the assemblage of primary producers and composition likely affected the survival of fish eggs and larvae through an impairment of resource acquisition (prevented negative effects) and avoidance of predators in the water column (presumed positive effects). Benthic infauna, including polychaetes, oligochaetes, and nematodes, respond positively to oil enrichment, which, for many fishes, likely increases available prey resources (D'Aleone et al. 1984; Truand and Burger 1994). However, it remains questionable whether benthic feeding fishes forage effectively within contaminated sediments (Cragg et al. 1997). While sub-marine and seagrass coverages, shrimp, crab larvae, and oysters are all highly sensitive to PAH

mechanisms that reduced the overall population impacts of the Macondo spill in sub-marine and seagrass habitats.

Benthic nekton. Many fishes are highly mobile and are likely capable of fleeing oil-affected shorelines, given the near and total gradients of disturbance in coastal habitats following the Macondo spill. For marine fishes, the ability to seek refugia confers resilience against the effects of hydrocarbon pollution associated with offshore petroleum production platforms, despite quantifiable impacts for immobile invertebrates (e.g., crustaceans, echinoderms, other invertebrate deposit feeders Peterson et al. 1985; Iskers, Fieldman and Schaeffer (e.g., spot, *Leiostomus xanthurus*) are capable of detecting and avoiding heavily oiled sediments, although they do not necessarily avoid lightly oiled sediments on food items (Miles et al. 1996; Heath-Crowe et al. 2001; <http://www.gomri.org/2010/09/06/2010-09-06-01>).

1998). Furthermore, long-term periodic exposures to hydrocarbons in regions with natural background seepage such as the northern GOM may prime adaptive avoidance behaviors or tolerance in resident species (sensu Rozas et al. 2000, Van Veld and Nacci 2008).

marsh and seagrass habitat loss associated with oiling may have negative effects on regional productivity over multiple generations (Mendelsohn et al. 2012), although such losses may not cause detectable site-specific decreases in fish density in the short term. Beyond habitat loss, oil contamination in sediments can remain detectable for decades in some habitats, with subtle, long-term effects on the fitness of sediment-associated species (Cahill et al. 2008).

Factors dampening population-level responses despite organismal ecotoxicity
Despite the significant oil and sulfidic threats posed by oil pollution, the life histories of many fishes in the GOM may have promoted avoidance behaviors or compensatory

coastal environments at concentrations sufficient to cause biological responses in resident fishes and that different cannot explain the divide between organismal and population-level findings.

Compensatory processes. Density-mediated increases in vital rates, such as juvenile and adult survival and growth rates, may often be sufficient to overcome the impacts of oil exposure, which may result in little change at the population level. Such compensatory responses are frequently quite strong, and they underlie the resilience of many fish populations to additional mortality sources, such as harvest (Dunbar et al. 2013). For example, a meta-analysis of stock-recruitment relationships (i.e., the survival rate between egg and juvenile

September 2011 / Vol. 6 (No. 9) • Reference 785

CONFIDENTIAL

BP-HZN-2179MDL09309263

TREX-013287.0008