

Acute Embryonic or Juvenile Exposure to Deepwater Horizon Crude Oil Impairs the Swimming Performance of Mahi-Mahi (*Coryphaena hippurus*)

Edward M. Mager,^{*,†} Andrew J. Esbaugh,[‡] John D. Stieglitz,[†] Ronald Hoenig,[†] Charlotte Bodinier,[†] John P. Incardona,[§] Nathaniel L. Scholz,[§] Daniel D. Benetti,[†] and Martin Grosell[†]


[†]Division of Marine Biology and Fisheries, University of Miami, Rosenstiel School of Marine and Atmospheric Science, 4600 Rickenbacker Causeway, Miami, Florida 33149, United States

[‡]Department of Marine Science, University of Texas, Marine Science Institute, 750 Channel View Drive, Port Aransas, Texas 78373, United States

[§]Ecotoxicology Program, Northwest Fisheries Science Center, National Oceanic and Atmospheric Administration, 2725 Montlake Boulevard E, Seattle, Washington 98112, United States

Supporting Information

ABSTRACT: The Deepwater Horizon incident likely resulted in exposure of commercially and ecologically important fish species to crude oil during the sensitive early life stages. We show that brief exposure of a water-accommodated fraction of oil from the spill to mahi-mahi as juveniles, or as embryos/larvae that were then raised for ~25 days to juveniles, reduces their swimming performance. These physiological deficits, likely attributable to polycyclic aromatic hydrocarbons (PAHs), occurred at environmentally realistic exposure concentrations. Specifically, a 48 h exposure of $1.2 \pm 0.6 \mu\text{g L}^{-1}$ ΣPAHs (geometric mean \pm SEM) to embryos/larvae that were then raised to juvenile stage or a 24 h exposure of $30 \pm 7 \mu\text{g L}^{-1}$ ΣPAHs (geometric mean \pm SEM) directly to juveniles resulted in 37% and 22% decreases in critical swimming velocities (U_{crit}), respectively. Oil-exposed larvae from the 48 h exposure showed a 4.5-fold increase in the incidence of pericardial and yolk sac edema relative to controls. However, this larval cardiotoxicity did not manifest in a reduced aerobic scope in the surviving juveniles. Instead, respirometric analyses point to a reduction in swimming efficiency as a potential alternative or contributing mechanism for the observed decreases in U_{crit} .



INTRODUCTION

The largest marine oil spill in U.S. history spanned 87 days during the spring and summer of 2010 during which the blown-out Deepwater Horizon (DWH) Macondo wellhead released approximately 4 million barrels (6×10^8 L) of crude oil into the northern Gulf of Mexico (GoM).^{1–3} The DWH incident oiled large portions of the pelagic zone, including the upper surface waters, and overlapped both spatially and temporally with the spawning of a number of commercially and ecologically important fish species, including yellowfin and bluefin tuna,^{4,5} mahi-mahi,^{6,7} and other large pelagics.^{8–10} An unusual aspect of the spill was that it originated at depth and under high energy, which facilitated the dissolution of toxic polycyclic aromatic hydrocarbons (PAHs) by increasing mixing and contact time with the water during ascent to the surface.^{11,12} Thus, PAH exposures likely occurred for adult fish breeding in the area and during the far more sensitive early life stages (ELS; i.e., embryos and larvae) of the developing offspring. Besides immediate impacts on survival, such PAH exposures, even if brief, may have imparted subtle developmental and physiological impairments with ensuing ramifications of ecological importance.¹³ Here, we provide novel information on sublethal exposures of

the PAHs found in this oil spill to a resident GoM fish (mahi-mahi).

Numerous studies have investigated the developmental toxicity of crude oil to fish embryos and larvae and also identified the constituent PAHs responsible for ELS impacts.^{14–28} These functional studies have identified cardiac development as a key process impaired by PAHs, particularly 3-ring (tricyclic) compounds, causing anatomical malformations and functional defects that likely diminish cardiac output in association with bradycardia and arrhythmia. Oil-exposed embryos that develop cardiac failure and jaw deformities are unlikely to survive past the yolk sac stage to become free swimming and feeding larvae.¹⁴ These findings contravened a longstanding paradigm wherein lower molecular weight and more volatile 2-ring PAHs drive crude oil toxicity through acute and nonspecific narcosis.¹³ Recent studies have also shown that the nervous system may be a target of oil toxicity in fish, as

Received: April 2, 2014

Revised: May 21, 2014

Accepted: May 23, 2014

Published: May 23, 2014


ACS Publications

© 2014 American Chemical Society

7053

dx.doi.org/10.1021/es501628k | Environ. Sci. Technol. 2014, 48, 7053–7061

CONFIDENTIAL



BP-HZN-2179MDL09304810

heavy fuel oil exposure in marine fish embryos led to abnormal projections of cranial and peripheral nerves.^{29,30} Although severe oil effects culminate in cardiac failure, as evident from the accumulation of fluid around the heart or yolk sac (i.e., pericardial or yolk sac edema),^{17,19} more subtle changes in cardiac morphogenesis can cause permanent shape changes that persist to adulthood.¹⁶ However, the physiological consequences of cardiac and neurological defects in fish that survive transient PAH exposures remain poorly understood.

Given that ecologically important behaviors such as foraging, predator avoidance, and migration are limited by the maximum swimming velocity that a fish can sustain for a prolonged period (U_{crit}), tests of swimming performance provide a relevant measure of physiological fitness for fish exposed to relatively low levels of PAHs derived from crude oil.^{31,32} A recent study showed that embryonic exposure of zebrafish to Alaska North Slope crude oil (ANSO) conferred reduced U_{crit} in adulthood along with changes in ventricular shape, suggesting impaired cardiac output may have limited swimming performance in these fish.¹⁶ Still, similar studies are lacking for pelagic species with high aerobic demand, including the large open ocean fish that spawned in the surface spill zone during the DWH incident. In addition to the unknown latent effects following embryonic exposure, it remains to be seen whether transient oil exposure to GoM pelagic species during later stages of development (e.g., juvenile) might also cause immediate impairment to swimming performance.

The recent addition of a successful mahi-mahi husbandry program at the University of Miami Experimental Hatchery (UMEH) presents a rare opportunity to assess the short- and long-term impacts that the DWH incident may have imparted to this commercially and recreationally important GoM species during any life stage and under controlled laboratory conditions. In the present study, we tested the hypotheses that transient exposure to DWH oil prepared as a high-energy water-accommodated fraction (HEWAF) would (1) cause direct impairment to the swimming performance of juvenile mahi-mahi and (2) manifest in latent impairment as juveniles when exposed ~25 days earlier during the embryonic/larval stage. The oil used for exposures was collected on July 29, 2010 from a barge hold receiving slick oil from various skimmer vessels (hereafter referred to as slick A). Additionally, we sought to further investigate the role of reduced cardiac output as a potential underlying cause limiting swimming performance by assessing aerobic scope from respirometry measurements collected during the swim trials.

MATERIALS AND METHODS

Experimental Animals. Mahi-mahi broodstock (*Coryphaena hippurus*) were captured off the coast of Miami, FL, using hook and line angling techniques. The fish were subsequently transferred to the University of Miami Experimental Hatchery (UMEH), where they were acclimated in 80 m³ fiberglass maturation tanks (typically 5–7 per tank) equipped with recirculating aquaculture systems for water quality and temperature control. All embryos used in the experiments described herein were collected within ~2–10 h following a volitional (noninduced) spawn using standard UMEH methods.³³ A prophylactic formalin (37% formaldehyde solution) treatment was administered to the embryos (100 ppm for 1 h), followed by a 0.5 h rinse with a minimum of 300% water volume in the treatment vessel using filtered, UV-sterilized seawater. A small sample of eggs was collected from

each spawn to microscopically assess fertilization rate and embryo quality. Spawns with low fertilization rate (<85%) or frequent morphological abnormalities (>5%) were not used.

Preparation of Water-Accommodated Fractions. The oil (referred to herein as slick A) used to prepare all HEWAFs was collected during the DWH spill on July 29, 2010 from the hold of barge number CTC02404, which was receiving slick oil from various skimmer vessels (sample ID CTC02404-02), and was subsequently transferred under chain of custody to the University of Miami. Each HEWAF was prepared on the day of use at a loading rate of 1 g of oil per liter of 1 μ m filtered, UV-sterilized seawater and mixed in a Waring CB15 blender (Torrington, CT) at low speed for 30 s. The mixture was immediately transferred to a glass separatory funnel, allowed to settle for 1 h, and the lower ~90% carefully drained and retained for subsequent use as 100% WAF (unfiltered) that was diluted for test exposures.

Embryo–Larval Exposures (48 h). Embryos from the same cohort were exposed to control seawater or a 0.2% HEWAF dilution (1.2 μ g L⁻¹ Σ PAH) for 48 h and then raised for ~25 days to the early juvenile stage in clean seawater. The 0.2% HEWAF dilution was selected to target a concentration that minimized acute mortality yet was high enough to induce cardiotoxicity based on results from a previous slick A HEWAF bioassay. Exposures were established in a 2500 L cylindrical fiberglass tank (1 per treatment) coated with standard marine gelcoat paint on the interior and fitted with a PVC stand pipe that was filled with filtered (5–10 μ m), UV-sterilized seawater. For the oil exposure, HEWAF was added when the tank was approximately one-third filled by even distribution along a concentric path midway between the standpipe and the tank wall to ensure adequate mixing. Once the tanks were filled (~2200 L final volumes) flow was stopped and ~30 000–80 000 viable mahi-mahi embryos were stocked into each at <12 h postfertilization (hpf). A low level of aeration was provided to each tank using an air stone to distribute the embryos throughout the tank during static exposure. Following the 48 h exposure period, inflow of clean water to both tanks resumed and each remained under flow-through conditions with gentle aeration for the remainder of the experiment. Larvae were reared using UMEH intensive marine fish larviculture protocols and collected for swim trials at 25–26 days posthatch (dph). Specifically, larvae were pulse-fed enriched rotifers (*Brachionus plicatilis*) and *Artemia* nauplii during the first 2–3 weeks posthatch, followed by weaning onto an inert diet (Otohime, Marubeni Nisshin Feed Co., Ltd.). Waste was removed daily from each tank by siphoning.

Juvenile Exposures (24 h). Juveniles raised entirely in clean seawater (mean age of 31 d) were exposed to control seawater or 0.4, 1.2, or 2% HEWAF dilutions (4.2, 17, or 30 μ g L⁻¹ Σ PAH, respectively) for 24 h. Exposures were performed in a temperature controlled environmental chamber by placing six fish into 10–12 L of control seawater or seawater spiked with freshly prepared HEWAF in a 20 L glass jar with light aeration provided by an air stone. Temperature and photoperiod within the chamber were set at 27 °C and 16:8 h of light:dark, respectively. Fish were transferred directly from treatment tanks to the swimming respirometers (i.e., no recovery period was permitted prior to transfer). All swim trials were performed with clean seawater. Although only four fish were swum per replicate, six fish were exposed to safeguard against losses to mortality. With the exception of two fish that died within one of the 30 μ g L⁻¹ Σ PAH replicates, no other mortalities were

observed for the 24 h exposures. In all other replicates, the remaining two fish not used in the swim trials were euthanized. In sum, the number of replicates (and total n) for the control, 4.2, 17, and 30 $\mu\text{g L}^{-1}$ ΣPAH 24 h exposures were 5 (20), 2 (8), 3 (12) and 3 (12), respectively. Due to the rapid growth rates of juvenile mahi-mahi,³⁴ control exposures were interspersed among oil exposures to minimize potential confounding factors due to size on swimming performance and aerobic scope. Fish were obtained from three cohorts, with at least one control replicate included from each. Fish were fed in the morning before transferring to an exposure chamber, but not fed during the 24 h exposure period.

Water Quality and PAH Analysis. In addition to samples collected for ΣPAH analysis, the following water quality parameters were monitored: temperature, pH, dissolved oxygen (DO), salinity, and total ammonia. Temperature and DO were measured using a ProODO hand held optical DO probe and meter (YSI, Inc., Yellow Springs, OH) and pH was measured using a PHM201 meter (Radiometer, Copenhagen, Denmark) fitted with a combination glass electrode. The pH and DO probes were calibrated daily. Salinity was measured using a refractometer and total ammonia determined using a colorimetric assay.³⁵ All samples for PAH analysis were collected directly from exposure tanks as grab samples in 250 mL amber bottles and shipped overnight on ice to ALS Environmental (Kelso, WA) for analysis by gas chromatography/mass spectrometry—selective ion monitoring (GC/MS—SIM; based on EPA method 8270D). Reported ΣPAH values represent the sum of 50 select PAH analytes (Supporting Information Table S1; Figure S1).

For the 48 h embryo–larval exposures, ΣPAH samples and measurements for all water quality parameters except ammonia (final only) were collected daily during the exposure period (i.e., 0, 24, and 48 h) and at 24 and 48 h of the ensuing washout period when flow was returned to the tanks. For the 24 h juvenile exposures, initial and final measurements were taken for all parameters except ammonia (final only) and initial and final samples were collected for ΣPAH analysis. A summary of all measured water quality parameters and ΣPAH concentrations is provided in Supporting Information Table S2.

Assessment of Cardiac Abnormalities. Immediately following the 48 h exposure period, a subsample of larvae was collected from each of the grow-out tanks for assessment of pericardial edema and heart rate. Larvae ($n = 20$ per tank) were mounted 2–3 at a time over 2% methylcellulose in seawater and imaged using a Fire-i400 digital camera (Unibrain, San Ramon, CA) mounted on a Nikon SMZ800 stereomicroscope. Images were collected on a MacBook laptop using iMovie software and calibrated using a stage micrometer. Larvae were scored blind for presence or absence of pericardial edema by morphological assessment of the yolk mass which became distorted with pericardial edema. Presence of edema was noted if fluid accumulation was sufficient to distort the normal smooth bullet shape of the anterior yolk mass. Edema typically occurred in the form of a concave or pointed wedge shape of the anterior yolk mass and/or contained a number of smaller indentations indicated by dark, angular lines. Pericardial area was measured using ImageJ version 1.46r (rsbweb.nih.gov/ij/) from a perimeter drawn with the freehand tool enclosing this area (Supporting Information Figure S2). Lines were drawn across the boundary of the yolk sac only if distortion of the yolk mass was clearly evident by a sharp dark line. Heart rate was

measured by counting the total number of heart beats for each larva in a ≥ 10 s digital video clip played back at half speed.

Swimming Performance and Aerobic Scope. Four miniature (0.17 L) Blazka-style swim respirometers (Loligo Systems, Denmark) were used to assess U_{crit} and aerobic scope via automated intermittent flow respirometry.^{36,37} Oxygen consumption within each swim chamber was measured using a Pt100 fiber-optic probe connected to a Fibox 3 minisensor oxygen meter (PreSens Precision Sensing GmbH, Germany). The oxygen sensor was calibrated daily using 100% air saturation, established by vigorous aeration with an air stone, and 0% O_2 saturation, achieved using a solution of 10 g L^{-1} Na_2SO_3 (Sigma-Aldrich, St. Louis, MO). Temperature was maintained at 27 ± 1 °C using an aquarium heater submerged in the reservoir water surrounding each swim chamber and was measured through the Fibox meter using a separate probe. All data were collected using AutoResp2 version 2.1.2 (Loligo Systems, Denmark).

A preliminary study was performed with four juvenile fish to determine a minimum habituation period sufficient to permit recovery from handling stress and obtain a stable routine metabolic rate (RMR) prior to initiating a swim trial. The fish were each held in a swim chamber for 20 h at a minimal flow speed to maintain water mixing within the chamber without forcing the fish to swim. From these experiments, a stabilized RMR was established for each fish by 3–4 h. Thus, all subsequent swim trials followed a 4 h habituation period.

To measure U_{crit} , fish were forced to swim for 20 min intervals until fatigued using fixed velocity increments that, depending on size, ranged between 1 and 2.5 cm s^{-1} (~ 0.5 body lengths (BL s^{-1}) per interval. Unlike many other teleosts, mahi-mahi rarely exhibit fatigue by becoming pinned parallel to the rear screen. Rather, juveniles tend to maintain a position perpendicular to the rear screen while resting on a bent caudal fin, sometimes with intermittent swim bursts off the screen. Therefore, fatigue was designated as when the fish began to consistently rest on its caudal fin for several seconds at a time. U_{crit} (in BL s^{-1}) was calculated using the following equation originally described by Brett:³⁸ $U_{\text{crit}} = [U_f + (T/t)dU]/\text{cm}$, where U_f (cm s^{-1}) is the highest swim velocity maintained for a full interval, T (s) is the time spent at the final velocity, t is the time interval (s), and dU is the increment in swim speed (cm s^{-1}). Fish mass and BL were determined postswimming to minimize stress pretesting and to convert U_{crit} to BL s^{-1} . Food was withheld for all fish for at least 24 h prior to introduction into a swim chamber to eliminate digestion (i.e., specific dynamic action) as a confounding factor.

Although juveniles from the embryonic grow-out exposures were collected at a younger age and thus were significantly smaller (Table 1; $P < 0.05$), there was no significant difference in U_{crit} among any of the control cohorts. Furthermore, given that the controls were represented by a total of six treatment replicates from four cohorts, there was no indication of tank or batch effects on swimming performance. Therefore, fish from all of the control treatments were pooled for statistical comparisons of U_{crit} . This represents a conservative approach since smaller fish typically have higher U_{crit} values than larger fish when expressed in terms of body length³⁹ and the oil-exposed juveniles from the grow-out exposure were of smaller size than the combined controls (Table 1; $P < 0.05$).

A standardized approach was used to estimate aerobic scope for each fish. First, the logarithm of oxygen consumption ($\text{mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$) versus swimming speed (BL s^{-1}) was plotted for

Table 1. Biometric Data for Mahi-Mahi Used in Swimming Performance Tests^a

treatment	n	mass (mg)	BL (cm)	age (dph)
24 h Juvenile				
control	20 (17)	645 ± 97	4.4 ± 0.2	32 ± 2
0.4% HEWAF	8 (7)	404 ± 33	4.0 ± 0.1	28 ± 0
1.2% HEWAF	12 (9)	497 ± 29	4.3 ± 0.1	30 ± 0
2% HEWAF	12 (8)	812 ± 180	4.8 ± 0.3	35 ± 2
48 hpf				
control	5 (4)	283 ± 29	3.4 ± 0.1	26 ± 0
0.2% HEWAF	5 (5)	204 ± 35	3.1 ± 0.1	25 ± 0

^aNumbers in parentheses are *n* used in respirometry calculations ($r^2 \geq 0.7$ for linear regression of log O_2 consumption vs swimming speed). Values represent mean ± SEM. Abbreviations: hpf, hours postfertilization; dph, days postfertilization.

each fish and a least-squares linear regression performed (see Supporting Information Figure S3 for example). Standard metabolic rate (SMR; *y* intercept) and maximum metabolic rate (MMR; extrapolated at U_{crit}) were then derived from the resulting equations. Only individuals yielding a regression with an $r^2 \geq 0.7$ were used for these analyses (see Table 1 for *n*). Given the variation in observed body masses (Table 1), and that metabolic rate is known to scale with mass, we normalized for the effect of mass in calculating SMR and MMR by scaling to a standard mass of 1 g before calculating aerobic scope (MMR – SMR; Supporting Information Figure S4).

Cost of transport at U_{crit} ($COT_{U_{crit}}$) was calculated by dividing the MMR ($mg\ O_2\ g^{-1}\ s^{-1}$) for each fish by the corresponding U_{crit} ($BL\ s^{-1}$).⁴⁰

Statistical Analyses. Data are presented as means ± 1 standard error of the mean (SEM). Differences were tested for statistical significance using SigmaStat version 3.5 (Systat Software, Inc., San Jose, CA). Unless otherwise noted, see figure legends and table footnotes for the exact statistical method used for each test. In all cases, differences were deemed significant at $P < 0.05$.

RESULTS

PAH Concentrations and Composition of HEWAF Preparations. Because ΣPAH concentrations decreased over time, both the initial and geometric mean concentrations are reported (Supporting Information Table S2). Previous work has demonstrated consistency between PAH profiles obtained from our HEWAF preparations and samples collected from the active spill zone.²⁷ The percent composition of PAHs was similar among all oil exposures, with the bicyclic naphthalenes and tricyclic fluorenes, dibenzothiophenes, and phenanthrenes/anthracenes predominating (Supporting Information Figure S1). Nominal doses of HEWAF produced nearly proportional changes in ΣPAHs, although this relationship appeared to weaken at lower doses (Supporting Information Figure S1). Initial C1-phenanthrene concentrations (a PAH class well represented in DWH crude oil; Supporting Information Figure S1) ranged from 4.6–0.33 $\mu g\ L^{-1}$ in the 2% and 0.2% HEWAF dilutions, respectively (Supporting Information Table S1). Notably, the latter concentration falls below a previously reported level of $\geq 0.36\ \mu g\ L^{-1}$ representing 4.6% (25 of 548) of samples collected within a ~96 000 km² region centered over the wellhead.²⁷

Evidence of 48 h Larval Cardiotoxicity. Larvae that were exposed to oil for 48 h during early development (1.2 $\mu g\ L^{-1}$

ΣPAHs) were microscopically examined for defects in heart form (increased pericardial area due to edema) and function (heart rate) immediately following exposure using blind scoring of keyed images and videos. The incidence of pericardial edema increased 4.5-fold in oil-exposed larvae relative to controls (Figure 1A). Pericardial area also increased nearly 2-fold in




Figure 1. Evidence of cardiac toxicity in larval mahi-mahi exposed to 1.2 $\mu g\ L^{-1}$ ΣPAHs (geometric mean) for 48 h initiated <12 hpf (*n* = 20 each, except *n* = 19 for the oil-exposed group in (B)) as assessed by (A) percent incidence of pericardial edema, (B) pericardial area and (C) heart rate. Data in (B) are presented as means ± SEM. Data in (C) are presented as box plots indicating the 25th and 75th percentiles; whiskers indicate the 90th and 10th percentiles; filled circles indicate outliers; solid and dashed lines indicate the median and mean, respectively. *Significantly different from controls by Mann–Whitney rank sum test.

larvae that were previously exposed to HEWAF (Figure 1B; $P = 0.003$). Analysis of heart rates revealed greater variability in larvae exposed to HEWAF (Figure 1C), although the mean heart rate was not significantly different from control (Student's *t*-test; $P = 0.128$). Gross malformations that typify the more severe oil injury syndrome in fish ELS, including spinal, craniofacial and fin deformities,¹⁹ were not observed.




Figure 2. Comparisons of (A) critical swimming speed (U_{crit}), (B) standard metabolic rate (SMR), (C) maximum metabolic rate (MMR), and (D) aerobic scope from swimming respirometry experiments of juvenile mahi-mahi exposed to 4.2, 17, or 30 $\mu\text{g L}^{-1}$ ΣPAHs ($n = 7, 9$, and 8, respectively) for 24 h prior to swimming (shaded bars) or to 1.2 $\mu\text{g L}^{-1}$ ΣPAHs ($n = 5$) for 48 h initiated <12 hpf (hatched bar) and then raised for ~25 days to the juvenile stage. The control group represents fish from all four control cohorts combined ($n = 21$). Data in (B–C) were normalized for body mass by scaling to the respective SMR and MMR values predicted for a 1 g fish (aerobic scope = normalized MMR – normalized SMR). Data are presented as means \pm SEM. *Significantly different from controls by one-way ANOVA and Holm-Sidak multiple comparison procedure.

Swimming Performance. Results from the swim trials revealed a significant 37% ($P < 0.05$) decrease in mean U_{crit} for juveniles that were transiently exposed to 1.2 $\mu\text{g L}^{-1}$ ΣPAH as embryos and larvae. By comparison, among mahi-mahi exposed as juveniles, only the highest concentration (30 $\mu\text{g L}^{-1}$ ΣPAH) elicited a significant decrease of 22% ($P < 0.05$) (Figure 2A).

Metabolic Rates, Aerobic Scope, and Cost of Transport. Respirometry was conducted during each swim trial to determine the potential impacts of slick A HEWAF exposure on aerobic scope (i.e., maximum metabolic rate (MMR) – standard metabolic rate (SMR)). To account for variation in metabolic rates due to variation in size, calculations of SMR and MMR were first normalized for mass using empirically determined scaling coefficients and constants (see Materials and Methods). Non-normalized data are provided in the Supporting Information (Figure S5).

No significant effect on SMR, MMR, or aerobic scope was evident for the fish exposed to 1.2 $\mu\text{g L}^{-1}$ ΣPAH as embryos/larvae and then tested as juveniles ~25 days later (Figure 2). Accordingly, cost of transport at U_{crit} ($COT_{U_{crit}}$), which represents the mass-specific amount of aerobic energy expended to traverse a given distance,⁴⁰ increased significantly by 40% ($P = 0.0085$) to a level nearly matching the 37% decrease in U_{crit} (Figure 3). Similarly, no significant effect on SMR, MMR or aerobic scope was evident for the fish exposed to HEWAF as juveniles (Figure 2). Again, however, there was an upward trend in $COT_{U_{crit}}$ (21%, $P = 0.147$) at the 30 $\mu\text{g L}^{-1}$ ΣPAH concentration that closely mirrored the 22% decrease in U_{crit} (Figure 3).

DISCUSSION

Results from the present study demonstrate that exposure to crude oil collected from surface slicks during the DWH incident impairs the swimming performance of juvenile mahi-mahi, an




Figure 3. Cost of transport at U_{crit} for juvenile mahi-mahi exposed to 4.2, 17, or 30 $\mu\text{g L}^{-1}$ ΣPAHs ($n = 7, 9$, and 8, respectively) for 24 h prior to swimming (shaded bars) or to 1.2 $\mu\text{g L}^{-1}$ ΣPAHs ($n = 5$) for 48 h initiated <12 hpf (hatched bar) and then raised for ~25 days to the juvenile stage. The control group represents fish from all four control cohorts combined ($n = 21$). Data are presented as means \pm SEM. *Significantly different from controls by one-tailed Student's t -test and Bonferroni multiple comparison procedure.

ecologically and commercially important pelagic predator in the northern GoM. Perhaps most significant was the finding that such impairment during the juvenile stage manifested as a latent effect ~25 days following a 48 h exposure during the embryonic/larval stage to a ΣPAH concentration of 1.2 $\mu\text{g L}^{-1}$. By contrast, a nearly 30-times higher, yet still environmentally relevant, concentration of 30 $\mu\text{g L}^{-1}$ ΣPAH was required to elicit a similar decrease in U_{crit} for fish exposed for 24 h as juveniles. Although these results are consistent with an expected decrease in sensitivity with increasing developmental stage, they also support the suggestion that PAH exposure

during the embryonic/larval stage caused latent developmental effects. The effective embryonic/larval exposure was 1/30th that for the juveniles and the larval body burden would be expected to decrease over time through growth dilution and hepatic metabolism, potentially eliminating an effect of a residual body burden of PAHs that persisted to the juvenile stage. The observed larval effects of pericardial and yolk sac edema, which were similar to those previously described for a number of different teleosts exposed to various sources of crude oil,^{14,18,20,25,27,41–43} lend further credence to a likely early developmental origin for the latent effects on swimming performance.

The available data to date indicate that our exposure concentrations, which ranged from 1.2–30 $\mu\text{g L}^{-1}$ ΣPAHs , likely represent environmentally realistic exposure scenarios. For example, Bejarano et al. reported a range of <0.01–77 $\mu\text{g L}^{-1}$ ΣPAHs from samples collected at 1 and 10 m depths⁴⁴ and other studies have reported similar values ranging as high as 59 $\mu\text{g L}^{-1}$ and 85 $\mu\text{g L}^{-1}$ ΣPAHs from samples collected both at depth and at the surface within the active spill zone.^{45,46} However, it should be noted that comparisons among ΣPAH measurements are somewhat complicated by analytical and sample variations (e.g., the number and selection of analytes included in ΣPAH calculations and collection depths and locations). Additional perspective can be gained by comparing measurements of a single analyte, C1-phenanthrene, which is well represented within DWH crude oil. It was previously found that 4.6% of 548 samples collected from a region centered above the wellhead met or exceeded a concentration of 0.36 $\mu\text{g L}^{-1}$ for C1-phenanthrene, a value near the initial concentration of 0.33 $\mu\text{g L}^{-1}$ measured for the 0.2% HEWAF embryonic/larval exposure.²⁷ The fertilized eggs of mahi-mahi float near the surface of the open waters of the northern GoM where they are spawned and putatively remain to develop throughout their ELS history.⁷ Thus, given that the spill likely overlapped temporally and spatially with mahi-mahi spawning events,^{67,10} it seems highly probable that detrimental PAH exposures occurred for these fish during the sensitive ELSs.

Considering the above, the current findings likely indicate an additional level of reduced survival to maturity in the wild than might be anticipated from acute mortality alone due to the spill. For example, the latent effects exhibited by juveniles exposed as embryos/larvae support a potential mechanism of impaired swimming performance for the delayed mortality and, by extension, population-scale impacts shown previously by mark-recapture studies of pink salmon exposed to ANSCO (~5–19 $\mu\text{g L}^{-1}$ initial ΣPAH concentrations).^{47,48} The more direct effects on fish exposed transiently as juveniles may contribute yet a further level of mortality due to impaired swimming performance; however, the long-term impact on survival in this scenario is less clear as the likely duration of such impairment remains unknown at this time. Nevertheless, it is important to note that the effective concentrations reported herein may underestimate toxicity in the natural environment where other stressors with the potential to increase toxicant sensitivity (e.g., hypoxia, temperature, and UV exposure) are commonly present. Furthermore, such effects on mahi-mahi likely also span to other important pelagic GoM species, such as billfish and tunas, which also resided in the spill path during the ELSs.^{4,5,9,10}

There are several physiological/behavioral mechanisms that could potentially contribute to a reduced U_{crit} . First, U_{crit} may decrease as result of diminished aerobic scope deriving from a

limitation in oxygen uptake or delivery that decreases MMR (e.g., reduced cardiac output, anemia, or increased lamellar diffusion distance) or an effect that adds to the basal metabolic load and thus increases SMR (e.g., specific dynamic action, metabolism of xenobiotics, or increased ionoregulatory costs).⁴⁹ Second, glycolytically fuelled white muscles may be recruited to sustain high swimming speeds approaching and including U_{crit} .⁵⁰ Thus, any impairment to this recruitment may also limit U_{crit} . Finally, there is evidence that behavior plays a role in setting swimming speed when fish cannot generate or sense a ground speed, a particular concern when using swim tunnels where forward movement is greatly limited.⁵⁰ Although each of these potential mechanisms could conceivably play a role in reducing the U_{crit} of oil-exposed fish in our experiments, we chose to focus our investigation on the first mechanism of diminished aerobic scope in light of the well-known cardiotoxic effects of PAHs.

A recent study revealed that zebrafish that survived to adulthood following a 48 h embryonic exposure to ANSCO (24–36 $\mu\text{g L}^{-1}$ ΣPAHs) developed defects in ventricular shape likely to impair cardiac output.¹⁶ Consistent with this, swim trials with the affected adult zebrafish revealed an 18% decrease in U_{crit} , thereby supporting a link between impaired cardiac output, reduced aerobic scope, and impaired swimming performance.¹⁶ However, respirometry was not performed in the zebrafish trials, and we did not examine the hearts of juvenile mahi-mahi for permanent changes in heart shape in response to embryonic PAH exposure. Nevertheless, larval evidence for developmental cardiotoxicity (pericardial and yolk sac edema) following the 48 h exposure period would suggest that subsequent reductions in juvenile swimming performance might be a consequence of a reduced aerobic scope, with a diminished cardiac output limiting MMR. Yet, our respirometry results did not clearly reveal a reduced MMR as the underlying mechanism for the impaired swimming ability of juvenile mahi-mahi. Although direct measurements of cardiac performance were not made, the lack of an effect on aerobic scope strongly indicates that cardiac output was maintained, thereby suggesting a mechanism(s) for the reduced U_{crit} other than, or in addition to, one that limits aerobic scope for juvenile mahi-mahi exposed to DWH crude oil.

The apparent discrepancy between potential aerobic scope effects observed in zebrafish but not juvenile mahi-mahi may relate to species-specific differences in life histories, most notably the aggressive predatory nature of the latter. Mahi-mahi exhibit a high degree of cannibalism when reared at high density under standard culture conditions. This was frequently observed during our rearing procedures and undoubtedly introduced a baseline level of selection that would have only been exacerbated with the addition of oil. Thus, the absence of a corresponding reduced aerobic scope by juvenile fish from the same cohort exhibiting larval cardiotoxicity may indicate that the most severely impaired fish with cardiac defects were selected out during the rearing process. Similar fish in the wild, however, would not have been subjected to the same high selective pressures due to confinement but rather pressures from other species and environmental stressors as discussed above. In the end, impaired juvenile swimming performance by feral fish exposed to crude oil may result both from an as of yet undetermined physiological deficit, as indicated by the present study, and a reduced aerobic scope owing to developmental cardiac defects as indicated by other studies. Nevertheless, considering that in all likelihood the most highly fit fish were

used for the swim trials, in addition to the fact that the exposures were performed in the absence of environmental stressors, lends further support that our results are likely conservative.

The observed decreases in U_{crit} without corresponding decreases in aerobic scope as described herein suggest that the former are due to reductions in swimming efficiency (i.e., greater energy is required to swim a given distance or maintain a given velocity). This is well illustrated by the proportional inverse relationships for U_{crit} and $COT_{U_{crit}}$ exhibited by both the juveniles swam ~25 days following exposure to HEWAF as embryos/larvae and the juveniles swam 24 h after exposure to the highest concentration of HEWAF. Thus, it is plausible that the latent and acute effects are related to direct toxic effects in the red or white swimming muscles, or the neural control of swimming muscle contractions, that cause dysfunction in the proper coordination of muscles required for efficient swimming. For example, this might occur due to a PAH-induced disruption in the excitation-contraction coupling mechanism of skeletal muscles in manner consistent with that recently described for bluefin and yellowfin tuna²⁶ cardiomyocytes. Alternatively, PAH-induced ROS production could result in the uncoupling of mitochondrial oxidative mechanisms that diminish the amount of ATP generated per unit oxygen.^{51,52} Considering that both sustained aerobic activity and anaerobic burst swimming are utilized in achieving U_{crit} , a failure in gait transition to anaerobic burst swimming by white muscles might represent another possible explanation as greater demand would be placed on aerobic metabolism to sustain U_{crit} .⁵³ Finally, an intact lateral line is necessary for efficient swimming⁵⁴ and an oil-induced abnormality in lateral line development could underlie the observed changes in cost of transport. Taken together, these considerations may implicate a role for neurological dysfunction in limiting the swimming performance of pelagic fish species exposed to DWH crude oil. Clearly, more research is needed to test these hypotheses and elucidate the underlying mechanism(s) of impaired swimming performance by juvenile mahi-mahi following transient exposure to DWH crude oil.

Irrespective of mechanism, our findings of reduced U_{crit} for mahi-mahi exposed to DWH crude oil nevertheless reveal significant physiological impairment at PAH concentrations well below some measurements taken during the spill (i.e., as low as $1.2 \mu\text{g L}^{-1}$ ΣPAHs). These results indicate that the impacts on survival of mahi-mahi to maturity are likely to be greater than estimates that consider acute mortality alone. Still, there remains much to be learned regarding the effects of DWH oil exposure on the swimming performance of resident GoM pelagic fishes, including for example the ability to recover from transient exposures and the significance of natural environmental stressors in contributing additional detriment. Such information would provide further insight relevant to assessing the damage that the DWH spill imparted to these important natural resources.

■ ASSOCIATED CONTENT

Supporting Information

Further information is available that provides PAH compositions and measurements (Tables S1 and S2; Figure S1), water quality measurements (Table S2), an example of larval pericardial edema (Figure S2), an example linear regression of metabolic rate vs swimming speed (Figure S3), methods for determining mass scaling constants and coefficients (Figure

S4), and nonmass scaled metabolic rate and COT data (Figure S5). This material is available free of charge via the Internet at <http://pubs.acs.org/>.

■ AUTHOR INFORMATION

Corresponding Author

*E. M. Mager. E-mail: emager@rsmas.miami.edu. Phone: 305-421-4823. Fax: 305-421-4600. Division of Marine Biology and Fisheries, University of Miami, Rosenstiel School of Marine and Atmospheric Science, 4600 Rickenbacker Cswy., Miami, FL 33149-1098, United States.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors would like to thank Dr. Anthony Farrell for his insightful review and suggestions for the manuscript, as well as Stratus Consulting for their valuable assistance and feedback, Kathleen Munley and Maria Rodgers for their technical support, and the students and staff at the UMEH for their assistance in the culture of mahi-mahi used in this study. This work was supported by funds provided as part of the Natural Resource Damage Assessment (NRDA) for the DWH oil spill. Data presented here are a subset of a larger toxicological database that is being generated as part of the DWH NRDA, therefore, these data will be subject to additional analysis and interpretation, which may include interpretation in the context of additional data not presented here.

■ REFERENCES

- (1) Camilli, R.; Di Iorio, D.; Bowen, A.; Reddy, C. M.; Techet, A. H.; Yoerger, D. R.; Whitcomb, L. L.; Seewald, J. S.; Sylva, S. P.; Fenwick, J. Acoustic measurement of the Deepwater Horizon Macondo well flow rate. *Proc. Natl. Acad. Sci. U. S. A.* 2012, 109 (50), 20235–20239.
- (2) Crone, T. J.; Tolstoy, M. Magnitude of the 2010 Gulf of Mexico oil leak. *Science* 2010, 330 (6004), 634.
- (3) McNutt, M.; Camilli, R.; Guthrie, G.; Hsieh, P.; Labson, V.; Lehr, B.; Maclay, D.; Ratzel, A.; Sogge, M. *Assessment of flow rate estimates for the Deepwater Horizon/Macondo well oil spill*; United States Department of the Interior: Washington, DC, 2011.
- (4) Block, B. A.; Teo, S. L.; Walli, A.; Boustany, A.; Stokesbury, M. J.; Farwell, C. J.; Weng, K. C.; Dewar, H.; Williams, T. D. Electronic tagging and population structure of Atlantic bluefin tuna. *Nature* 2005, 434 (7037), 1121–1127.
- (5) Muhling, B. A.; Roffer, M. A.; Lamkin, J. T.; Ingram, G. W., Jr.; Upton, M. A.; Gawlikowski, G.; Muller-Karger, F.; Habtes, S.; Richards, W. J. Overlap between Atlantic bluefin tuna spawning grounds and observed Deepwater Horizon surface oil in the northern Gulf of Mexico. *Mar. Pollut. Bull.* 2012, 64 (4), 679–687.
- (6) Gibbs, R. H.; Collette, B. B. On the identification, distribution, and biology of the dolphins, *Coryphaena hippurus* and *C. equiselis*. *Bull. Mar. Sci.* 1959, 9, 117–152.
- (7) Palko, B. J.; Beardsley, G. L.; Richards, W. J. *Synopsis of the biological data on dolphin-fishes, Coryphaena hippurus Linnaeus and Coryphaena equiselis Linnaeus*; United States Department of Commerce: Washington, DC, 1982.
- (8) Brown-Peterson, N.; Overstreet, R. M.; Lotz, J. M.; Franks, J. S.; Burns, K. M. Reproductive biology of cobia, *Rachycentron canadum*, from coastal waters of the southern United States. *Fish. Bull.* 2001, 99, 15–28.
- (9) Rooker, J. R.; Kitchens, L. L.; Dance, M. A.; Wells, R. J.; Falterman, B.; Cornic, M. Spatial, temporal, and habitat-related variation in abundance of pelagic fishes in the Gulf of Mexico: Potential implications of the Deepwater Horizon oil spill. *PLoS One* 2013, 8 (10), e76080.

- (10) Rooker, J. R.; Simms, J. R.; Wells, R. J.; Holt, S. A.; Holt, G. J.; Graves, J. E.; Purey, N. B. Distribution and habitat associations of billfish and swordfish larvae across mesoscale features in the Gulf of Mexico. *PLoS One* 2012, 7 (4), e34180.
- (11) Reddy, C. M.; Arey, J. S.; Seewald, J. S.; Sylva, S. P.; Lemkau, K. L.; Nelson, R. K.; Carmichael, C. A.; McIntyre, C. P.; Fenwick, J.; Ventura, G. T.; Van Mooy, B. A.; Camilli, R. Composition and fate of gas and oil released to the water column during the Deepwater Horizon oil spill. *Proc. Natl. Acad. Sci. U. S. A.* 2012, 109 (50), 20229–20234.
- (12) Ryerson, T. B.; Camilli, R.; Kessler, J. D.; Kujawinski, E. B.; Reddy, C. M.; Valentine, D. L.; Atlas, E.; Blake, D. R.; de Gouw, J.; Meinardi, S.; Parrish, D. D.; Peischl, J.; Seewald, J. S.; Warneke, C. Chemical data quantify Deepwater Horizon hydrocarbon flow rate and environmental distribution. *Proc. Natl. Acad. Sci. U. S. A.* 2012, 109 (50), 20246–20253.
- (13) Peterson, C. H.; Rice, S. D.; Short, J. W.; Esler, D.; Bodkin, J. L.; Ballachey, B. E.; Irons, D. B. Long-term ecosystem response to the Exxon Valdez oil spill. *Science* 2003, 302 (5653), 2082–2086.
- (14) Carls, M. G.; Rice, S. D.; Hose, J. E. Sensitivity of fish embryos to weathered crude oil: Part I. Low-level exposure during incubation causes malformations, genetic damage, and mortality in larval Pacific herring (*Clupea pallasii*). *Environ. Toxicol. Chem.* 1999, 18, 481–493.
- (15) Fallahatafti, S.; Rantanen, T.; Brown, R. S.; Snieckus, V.; Hodson, P. V. Toxicity of hydroxylated alkyl-phenanthrenes to the early life stages of Japanese medaka (*Oryzias latipes*). *Aquat. Toxicol.* 2012, 106–107, 56–64.
- (16) Hicken, C. E.; Linbo, T. L.; Baldwin, D. H.; Willis, M. L.; Myers, M. S.; Holland, L.; Larsen, M. G.; Stekol, M. S.; Rice, S. D.; Collier, T. K.; Scholz, N. L.; Incardona, J. P. Sublethal exposure to crude oil during embryonic development alters cardiac morphology and reduces aerobic capacity in adult fish. *Proc. Natl. Acad. Sci. U. S. A.* 2011, 108 (17), 7086–7090.
- (17) Hodson, P. V.; Qureshi, K.; Noble, C. A.; Akhtar, P.; Brown, R. S. Inhibition of CYP1A enzymes by alpha-naphthoflavone causes both synergism and antagonism of retene toxicity to rainbow trout (*Oncorhynchus mykiss*). *Aquat. Toxicol.* 2007, 81 (3), 275–285.
- (18) Incardona, J. P.; Carls, M. G.; Day, H. L.; Sloan, C. A.; Bolton, J. L.; Collier, T. K.; Scholz, N. L. Cardiac arrhythmia is the primary response of embryonic Pacific herring (*Clupea pallasii*) exposed to crude oil during weathering. *Environ. Sci. Technol.* 2009, 43 (1), 201–207.
- (19) Incardona, J. P.; Carls, M. G.; Teraoka, H.; Sloan, C. A.; Collier, T. K.; Scholz, N. L. Aryl hydrocarbon receptor-independent toxicity of weathered crude oil during fish development. *Environ. Health Perspect.* 2005, 113 (12), 1755–1762.
- (20) Incardona, J. P.; Collier, T. K.; Scholz, N. L. Defects in cardiac function precede morphological abnormalities in fish embryos exposed to polycyclic aromatic hydrocarbons. *Toxicol. Appl. Pharmacol.* 2004, 196 (2), 191–205.
- (21) Incardona, J. P.; Day, H. L.; Collier, T. K.; Scholz, N. L. Developmental toxicity of 4-ring polycyclic aromatic hydrocarbons in zebrafish is differentially dependent on AH receptor isoforms and hepatic cytochrome P4501A metabolism. *Toxicol. Appl. Pharmacol.* 2006, 217 (3), 308–321.
- (22) Incardona, J. P.; Linbo, T. L.; Scholz, N. L. Cardiac toxicity of 5-ring polycyclic aromatic hydrocarbons is differentially dependent on the aryl hydrocarbon receptor 2 isoform during zebrafish development. *Toxicol. Appl. Pharmacol.* 2011, 257 (2), 242–249.
- (23) Marty, G. D.; Hinton, D. E.; Short, J. W.; Heintz, R. A.; Rice, S. D.; Dambach, D. M.; Willits, N. H.; Stegeman, J. J. Ascites, premature emergence, increased gonadal cell apoptosis, and cytochrome P4501A induction in pink salmon larvae continuously exposed to oil-contaminated gravel during development. *Can. J. Zool.* 1997, 75, 989–1007.
- (24) Scott, J. A.; Incardona, J. P.; Pelkki, K.; Shepardon, S.; Hodson, P. V. AhR2-mediated, CYP1A-independent cardiovascular toxicity in zebrafish (*Danio rerio*) embryos exposed to retene. *Aquat. Toxicol.* 2011, 101 (1), 165–174.
- (25) Turcotte, D.; Akhtar, P.; Bowerman, M.; Kiparissis, Y.; Brown, R. S.; Hodson, P. V. Measuring the toxicity of alkyl-phenanthrenes to early life stages of medaka (*Oryzias latipes*) using partition-controlled delivery. *Environ. Toxicol. Chem.* 2011, 30 (2), 487–495.
- (26) Brette, F.; Machado, B.; Cros, C.; Incardona, J. P.; Scholz, N. L.; Block, B. A. Crude oil impairs cardiac excitation-contraction coupling in fish. *Science* 2014, 343 (6172), 772–776.
- (27) Incardona, J. P.; Gardner, L. D.; Linbo, T. L.; Brown, T. L.; Esbaugh, A. J.; Mager, E. M.; Stieglitz, J. D.; French, B. L.; Labenia, J. S.; Lactz, C. A.; Tagal, M.; Sloan, C. A.; Elizur, A.; Benetti, D. D.; Grosell, M.; Block, B. A.; Scholz, N. L. Deepwater Horizon crude oil impacts the developing hearts of large predatory pelagic fish. *Proc. Natl. Acad. Sci. U. S. A.* 2014, No. 10.1073/pnas.1320950111.
- (28) de Soysa, T. Y.; Ulrich, A.; Friedrich, T.; Pite, D.; Compton, S. L.; Ok, D.; Bernardos, R. L.; Downes, G. B.; Hsieh, S.; Stein, R.; Lagdameo, M. C.; Halvorsen, K.; Kesich, L. R.; Barresi, M. J. Macondo crude oil from the Deepwater Horizon oil spill disrupts specific developmental processes during zebrafish embryogenesis. *BMC Biol.* 2012, 10, 40.
- (29) Irie, K.; Kawaguchi, M.; Mizuno, K.; Song, J. Y.; Nakayama, K.; Kitamura, S.; Murakami, Y. Effect of heavy oil on the development of the nervous system of floating and sinking teleost eggs. *Mar. Pollut. Bull.* 2011, 63 (5–12), 297–302.
- (30) Kawaguchi, M.; Song, J. Y.; Irie, K.; Murakami, Y.; Nakayama, K.; Kitamura, S. Disruption of Sema3A expression causes abnormal neural projection in heavy oil exposed Japanese flounder larvae. *Mar. Pollut. Bull.* 2011, 63 (5–12), 356–361.
- (31) Fuiman, L. A.; Rose, K. A.; Cowan, J. H.; Smith, E. P. Survival skills required for predator evasion by fish larvae and their relation to laboratory measures of performance. *Anim. Behav.* 2006, 71, 1389–1399.
- (32) Plaut, I. Critical swimming speed: its ecological relevance. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* 2001, 131 (1), 41–50.
- (33) Stieglitz, J. D.; Benetti, D. D.; Hoenig, R. H.; Sardenberg, B.; Welch, A. W.; Miralao, S. Environmentally conditioned, year-round volitional spawning of cobia (*Rachycentron canadum*) in broodstock maturation systems. *Aquacult. Res.* 2012, 43, 1557–1566.
- (34) Benetti, D. D.; Iversen, E. S.; Ostrowski, A. C. Growth rates of captive dolphin, *Coryphaena hippurus*, in Hawaii. *Fish. Bull.* 1995, 93, 152–157.
- (35) Verdouw, H.; van Ehteld, C. J. A.; Dekkers, F. M. J. Ammonia determination based on indophenol formation with sodium salicylate. *Water Res.* 1978, 12, 399–402.
- (36) Blazka, P.; Volf, M.; Cepela, M. A new type of respirometer for the determination of the metabolism of fish in an active state. *Physiol. Bohemoslov.* 1960, 9, 553–558.
- (37) Steffenson, J. F. Some errors in respirometry of aquatic breathers: how to avoid and correct for them. *Fish. Physiol. Biochem.* 1989, 6, 49–59.
- (38) Brett, J. R. The respiratory metabolism and swimming performance of young sockeye salmon. *J. Fish. Res. Board Can.* 1964, 21, 1183–1226.
- (39) Beamish, F. W. H. Swimming Capacity. In *Fish Physiology*; Hoar, W. S., Randall, D. J., Eds.; Academic Press: New York, 1978; Vol. VII, pp 101–187.
- (40) Videler, J. J. *Fish swimming*; Chapman & Hall: New York, 1993.
- (41) Couillard, C. M. A microscale test to measure petroleum oil toxicity to mummichog embryos. *Environ. Toxicol.* 2002, 17 (3), 195–202.
- (42) Jung, J. H.; Hicken, C. E.; Boyd, D.; Anulacion, B. F.; Carls, M. G.; Shim, W. J.; Incardona, J. P. Geologically distinct crude oils cause a common cardiotoxicity syndrome in developing zebrafish. *Chemosphere* 2012, 91 (8), 1146–1155.
- (43) Pollino, C. A.; Holdway, D. A. Toxicity testing of crude oil and related compounds using early life stages of the crimson-spotted rainbowfish (*Melanoaenia fluviatilis*). *Ecotoxicol. Environ. Saf.* 2002, 52, 180–189.
- (44) Bejarano, A. C.; Levine, E.; Mearns, A. J. Effectiveness and potential ecological effects of offshore surface dispersant use during

the Deepwater Horizon oil spill: a retrospective analysis of monitoring data. *Environ. Monit. Assess.* 2013, 185, 10281–10295.

(45) Diercks, A.-R.; Highsmith, R. C.; Asper, V. L.; Joung, D.; Zhou, Z.; Guo, L.; Shiller, A. M.; Joye, S. B.; Teske, A. P.; Guinasso, N.; Wade, T. L.; Lohrenz, S. E. Characterization of subsurface polycyclic aromatic hydrocarbons at the Deepwater Horizon site. *Geophys. Res. Lett.* 2010, 37, L20602.

(46) Wade, T. L.; Sweet, S. T.; Sericano, J. L.; Guinasso, N. L.; Diercks, A.-R.; Highsmith, R. C.; Asper, V. L.; Joung, D.; Shiller, A. M.; Lohrenz, S. E.; Joye, S. B. *Analyses of Water Samples From the Deepwater Horizon Oil Spill: Documentation of the Subsurface Plume*; American Geophysical Union: Washington DC, 2011; Vol. 195, p 77–82.

(47) Heintz, R. A. Chronic exposure to polynuclear aromatic hydrocarbons in natal habitats leads to decreased equilibrium size, growth, and stability of pink salmon populations. *Integr. Environ. Assess. Manage.* 2007, 3 (3), 351–363.

(48) Heintz, R. A.; Rice, S. D.; Wertheimer, A. C.; Bradshaw, R. F.; Thrower, F. P.; Joyce, J. E.; Short, J. W. Delayed effects on growth and marine survival of pink salmon *Oncorhynchus gorbuscha* after exposure to crude oil during embryonic development. *Mar. Ecol.: Prog. Ser.* 2000, 208, 205–216.

(49) Brett, J. R., Implications and assessments of environmental stress. In *The Investigation of Fish-Power Problems*; Institute of Fisheries, University of British Columbia: Vancouver, BC, 1958.

(50) Farrell, A. P. Cardiorespiratory performance during prolonged swimming tests with salmonids: a perspective on temperature effects and potential analytical pitfalls. *Philos. Trans. R. Soc., B* 2007, 362, 2017–2030.

(51) Jastroch, M.; Divakaruni, A. S.; Mukerjee, S.; Treberg, J. R.; Brand, M. D. Mitochondrial proton and electron leaks. *Essays Biochem.* 2010, 47, 53–67.

(52) Genova, M. L.; Lenaz, G. Functional role of mitochondrial respiratory supercomplexes. *Biochim. Biophys. Acta* 2014, 1837 (4), 427–443.

(53) Lee, C. G.; Farrell, A. P.; Lotto, A.; Hinch, S. G.; Healey, M. C. Excess post-exercise oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*) salmon following critical speed swimming. *J. Exp. Biol.* 2003, 206, 3253–3260.

(54) Yanase, K.; Herbert, N. A.; Montgomery, J. C. Disrupted flow sensing impairs hydrodynamic performance and increases the metabolic cost of swimming in the yellowtail kingfish, *Seriola lalandi*. *J. Exp. Biol.* 2012, 215, 3944–3954.

Table S1. Individual measurements ($\mu\text{g L}^{-1}$) for 50 (shaded) of 76 total analytes used in calculating ΣPAH values provided in Table S2 and percent compositions provided in Figure S1. Values represent the initial sample (0.2% HEWAF growout) or mean of initial samples (all others).

	2% HEWAF			1.2% HEWAF			0.4% HEWAF			0.2% HEWAF		
	Mean	% $\Sigma\text{Tot.}$	% $\Sigma 50$	Mean	% $\Sigma\text{Tot.}$	% $\Sigma 50$	Mean	% $\Sigma\text{Tot.}$	% $\Sigma 50$	Initial	% $\Sigma\text{Tot.}$	% $\Sigma 50$
1-Methyldibenzothiophene	0.17	0.28		0.10	0.28		0.03	0.30		0.01	0.43	
1-Methylnaphthalene	0.06	0.09		0.03	0.10		0.00	0.00		0.00	0.00	
1-Methylphenanthrene	0.97	1.63		0.56	1.64		0.17	1.62		0.07	2.34	
2,3,5-Trimethylnaphthalene	1.28	2.17		0.83	2.40		0.24	2.37		0.11	3.57	
2,6-Dimethylnaphthalene	0.62	1.05		0.35	1.02		0.13	1.23		0.04	1.42	
2/3-Methyldibenzothiophene	0.24	0.41		0.13	0.38		0.04	0.44		0.02	0.55	
2-Methylantracene	0.00	0.00		0.00	0.00		0.00	0.00		0.00	0.00	
2-Methylnaphthalene	0.05	0.08		0.02	0.07		0.00	0.00		0.00	0.00	
2-Methylphenanthrene	1.14	1.92		0.68	1.98		0.20	1.92		0.08	2.66	
3-Methylphenanthrene	1.01	1.70		0.56	1.61		0.17	1.67		0.07	2.19	
4/9-Methylphenanthrene	1.26	2.13		0.72	2.08		0.22	2.18		0.08	2.78	
4-Methyldibenzothiophene	0.53	0.89		0.31	0.90		0.09	0.90		0.05	1.48	
Acenaphthene	0.02	0.04	0.05	0.00	0.00	0.00	0.01	0.06	0.07	0.00	0.00	0.00
Acenaphthylene	0.00	0.00	0.00	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Anthracene	0.03	0.06	0.06	0.02	0.05	0.05	0.02	0.15	0.18	0.00	0.00	0.00
Benz(a)anthracene	0.04	0.07	0.09	0.03	0.08	0.09	0.01	0.13	0.16	0.01	0.45	0.55
Benzo(a)fluoranthene	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.16	0.20	0.00	0.00	0.00
Benzo(a)pyrene	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Benzo(b)fluoranthene	0.05	0.08	0.09	0.03	0.08	0.09	0.00	0.04	0.05	0.00	0.00	0.00
Benzo(b)fluorene	0.12	0.21	0.24	0.08	0.22	0.26	0.03	0.24	0.29	0.00	0.00	0.00
Benzo(b)thiophene	0.00	0.00		0.00	0.00		0.00	0.00		0.00	0.00	
Benzo(c)pyrene	0.09	0.15	0.17	0.05	0.14	0.17	0.02	0.16	0.19	0.00	0.00	0.00
Benzo(g,h,i)perylene	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.02	0.02	0.00	0.00	0.00
Benzo(j+k)fluoranthene	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Biphenyl	0.06	0.10	0.11	0.04	0.12	0.14	0.01	0.14	0.17	0.00	0.00	0.00
C1 - Chrysenes	0.86	1.46	1.72	0.52	1.50	1.75	0.16	1.58	1.89	0.07	2.45	2.97
C1 - Dibenzothiophenes	1.14	1.92	2.26	0.64	1.85	2.16	0.18	1.78	2.13	0.09	3.03	3.67

CONFIDENTIAL

BP-HZN-2179MDL09304820

C1 - Fluoranthenes/Pyrenes	0.62	1.05	1.24	0.33	0.96	1.12	0.11	1.05	1.25	0.00	0.00	0.00
C1 - Fluorenes	1.50	2.53	2.98	0.92	2.67	3.11	0.29	2.78	3.33	0.13	4.13	5.00
C1 - Benzothiophenes	0.00	0.00		0.00	0.00		0.02	0.15		0.00	0.00	
C1 - Decalins	0.00	0.00		0.00	0.00		0.02	0.17		0.00	0.00	
C1 - Naphthalenes	0.07	0.12	0.15	0.05	0.13	0.16	0.00	0.00	0.00	0.00	0.00	0.00
C1 - Naphthobenzothiophenes	0.59	0.99	1.16	0.34	0.99	1.15	0.12	1.16	1.39	0.00	0.00	0.00
C1 - Phenanthrenes/Anthracenes	4.57	7.71	9.08	2.72	7.88	9.20	0.79	7.63	9.15	0.33	10.92	13.23
C2 - Chrysenes	1.06	1.79	2.11	0.60	1.74	2.03	0.19	1.82	2.18	0.00	0.00	0.00
C2 - Dibenzo thiophenes	1.88	3.18	3.75	1.06	3.08	3.60	0.32	3.12	3.74	0.13	4.39	5.32
C2 - Fluoranthenes/Pyrenes	0.95	1.60	1.88	0.56	1.62	1.89	0.17	1.67	2.00	0.00	0.00	0.00
C2 - Fluorenes	2.73	4.61	5.43	1.68	4.89	5.70	0.49	4.73	5.67	0.19	6.20	7.51
C2 - Benzothiophenes	0.00	0.00		0.00	0.00		0.01	0.11		0.00	0.00	
C2 - Decalins	0.00	0.00		0.00	0.00		0.00	0.00		0.00	0.00	
C2 - Naphthalenes	1.71	2.89	3.41	1.07	3.10	3.62	0.31	3.01	3.60	0.16	5.08	6.16
C2 - Naphthobenzothiophenes	0.76	1.28	1.50	0.48	1.40	1.64	0.14	1.38	1.65	0.00	0.00	0.00
C2 - Phenanthrenes/Anthracenes	5.64	9.53	11.22	3.46	10.05	11.73	1.02	9.89	11.85	0.34	11.09	13.43
C3 - Chrysenes	0.70	1.18	1.39	0.34	0.98	1.15	0.14	1.36	1.63	0.00	0.00	0.00
C3 - Dibenzo thiophenes	1.43	2.42	2.85	0.88	2.56	2.98	0.24	2.31	2.77	0.12	3.90	4.73
C3 - Fluoranthenes/Pyrenes	1.20	2.03	2.39	0.64	1.87	2.18	0.20	1.92	2.30	0.00	0.00	0.00
C3 - Fluorenes	2.69	4.54	5.35	1.71	4.95	5.78	0.52	5.01	6.00	0.00	0.00	0.00
C30-Hopane	0.67	1.13		0.36	1.05		0.15	1.42		0.00	0.00	
C3 - Benzothiophenes	0.06	0.09		0.03	0.08		0.02	0.20		0.00	0.00	
C3 - Decalins	0.00	0.00		0.00	0.00		0.00	0.00		0.00	0.00	
C3 - Naphthalenes	4.48	7.56	8.90	2.78	8.06	9.40	0.64	6.18	7.41	0.28	9.15	11.08
C3 - Naphthobenzothiophenes	0.50	0.84	0.99	0.00	0.00	0.00	0.12	1.21	1.45	0.00	0.00	0.00
C3 - Phenanthrenes/Anthracenes	3.49	5.90	6.95	2.10	6.10	7.12	0.56	5.43	6.51	0.24	8.00	9.69
C4 - Chrysenes	0.23	0.39	0.46	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
C4 - Dibenzo thiophenes	1.72	2.90	3.41	1.40	4.06	4.74	0.08	0.80	0.96	0.00	0.00	0.00
C4 - Benzothiophenes	0.06	0.10		0.00	0.00		0.01	0.08		0.00	0.00	
C4 - Decalins	0.66	1.12		0.15	0.44		0.13	1.28		0.00	0.00	
C4 - Fluoranthenes/Pyrenes	0.96	1.63	1.92	0.19	0.56	0.65	0.15	1.41	1.69	0.00	0.00	0.00
C4 - Naphthalenes	3.24	5.47	6.44	2.03	5.89	6.87	0.53	5.11	6.12	0.22	7.22	8.74
C4 - Naphthobenzothiophenes	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
C4 - Phenanthrenes/Anthracenes	2.25	3.79	4.47	1.18	3.42	4.00	0.41	3.96	4.74	0.00	0.00	0.00
Carbazole	0.00	0.00		0.00	0.00		0.00	0.00		0.00	0.00	
Chrysene+Triphenylene	0.41	0.69	0.81	0.24	0.70	0.81	0.09	0.84	1.01	0.03	0.86	1.04

TREX-013338.000011

CONFIDENTIAL

cis/trans-Decalin	0.00	0.00		0.00	0.00		0.01	0.12		0.00	0.00	
Dibenz(a,h)anthracene	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Dibenzofuran	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.13	0.15	0.00	0.00	0.00
Dibenzothiophene	0.23	0.40	0.47	0.13	0.38	0.45	0.05	0.46	0.55	0.02	0.55	0.66
Fluoranthene	0.04	0.06	0.07	0.01	0.04	0.05	0.00	0.03	0.04	0.00	0.00	0.00
Fluorene	0.36	0.60	0.71	0.20	0.58	0.68	0.09	0.84	1.01	0.03	0.97	1.18
Indeno(1,2,3-cd)pyrene	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Naphthalene	0.04	0.07	0.09	0.01	0.02	0.02	0.05	0.44	0.52	0.00	0.00	0.00
Naphthobenzothiophene	0.14	0.23	0.28	0.08	0.22	0.26	0.03	0.32	0.38	0.00	0.00	0.00
Perylene	0.00	0.00		0.00	0.00		0.00	0.00		0.00	0.00	
Phenanthrene	1.56	2.63	3.10	0.87	2.53	2.96	0.29	2.83	3.39	0.13	4.17	5.04
Pyrene	0.15	0.25	0.29	0.08	0.22	0.26	0.02	0.24	0.28	0.00	0.00	0.00
Retene	0.14	0.23		0.09	0.26		0.03	0.32		0.00	0.00	
ΣTotal	59.21	100		34.46	100		10.30	100		3.05	100	
Σ50	50.31		100	29.53		100	8.60		100	2.52		100

BP-HZN-2179MDL09304821

Table S2. Water quality (mean \pm SEM) and Σ PAH data for slick A HEWAF exposures. Σ PAH concentrations were calculated in 2 ways: (1) using all samples collected daily from each replicate (geometric mean \pm SEM); (2) in brackets, using only initial samples from each replicate (mean \pm SEM). Sample sizes are provided in parentheses. All other water quality values reflect measurements collected daily during exposure period, except total ammonia (final only).

Treatment	Σ PAH ($\mu\text{g L}^{-1}$)	Water temp. (C)	pH	D.O. (mg L^{-1})	Salinity (ppt)	Tot. amm. (μM)
<i>24 h Juvenile</i>						
Control	0.05 \pm 0.04 (10) [0.04 \pm 0.02 (5)]	26.8 \pm 0.3	7.89 \pm 0.05	5.92 \pm 0.13	35 \pm 1	15.2 \pm 4.5
0.4% HEWAF	4.2 \pm 1.9 (4) [8.6 \pm 0.9 (2)]	28.0 \pm 0.7	7.99 \pm 0.11	5.89 \pm 0.27	33 \pm 1	5.1 \pm 2.9
1.2% HEWAF	17 \pm 5 (6) [30 \pm 1 (3)]	27.0 \pm 0	7.97 \pm 0.18	5.57 \pm 0.38	35 \pm 1	11.8 \pm 1.9
2% HEWAF	30 \pm 7 (6) [50 \pm 1 (3)]	26.3 \pm 0.3	7.94 \pm 0.08	6.00 \pm 0.29	36 \pm 0	24.2 \pm 4.2
<i>48 hpf</i>						
Control	0.04 \pm 0.01 (3) [0.06 (1)]	28.6 \pm 0.6	8.14 \pm 0.03	7.32 \pm 0.26	32 \pm 1	0
0.2% HEWAF*	1.2 \pm 0.6 (3) [2.5 (1)]	28.3 \pm 0.7	8.20 \pm 0.05	6.53 \pm 0.12	31 \pm 0	0

* Σ PAH concentrations for 24 h and 48 h post-exposure washout periods were 0.3 and 0.2 $\mu\text{g L}^{-1}$, respectively.

Abbreviations: hpf, hours post-fertilization




Figure S1. Initial (0.2% HEWAF growout) or mean initial (all others) percent composition for 50 PAH analytes as determined by GC-MS for each of the slick A HEWAF exposures.

Abbreviations: naphthalenes (N), biphenyl (BPH), acenaphthene (ACE), acenaphthylene (ACY), fluorenes (F), dibenzothiophenes (D), anthracene (ANT), phenanthrene (P), phenanthrenes/anthracenes (PA), chrysene + triphenylene (CT), chrysenes (C), naphthobenzothiophenes (NB), fluoranthenes (FA), benz(a)anthracene (BAA), benzo(a)fluoranthene (BAF), benzo(a)pyrene (BAP), benzo(b)fluoranthene (BBF), benzo(b)fluorene (BF), benzo(e)pyrene (BEP), benzo(g,h,i)perylene (BGP), benzo(j+k)fluoranthene (BJF), dibenz(a,h)anthracene (DBA), dibenzofuran (DBF), indeno(1,2,3-cd)pyrene (IND) and pyrene (PYR). Numbers indicate quantity of carbons added per alkylated homolog.

CONFIDENTIAL

BP-HZN-2179MDL09304824

TREX-013338.000015




Figure S2. Representative still frame captures of larval mahi-mahi following 48 h exposure to (A) control conditions or (B) $1.2 \mu\text{g L}^{-1} \Sigma\text{PAHs}$ for 48 h initiated <12 hpf. Pericardial edema was measured as the area within a perimeter drawn from a freehand trace in ImageJ following the borders of the anterior pericardial region and contours of the yolk mass along the left side of the larvae. Example traces are illustrated by red dashed lines.




Figure S3. Example least-squares linear regression of metabolic rate vs. swimming speed (U) obtained from an individual juvenile mahi-mahi used in the swimming respirometry trials. Regressions yielding an $r^2 \geq 0.7$ were used for calculating SMR (y-intercept), MMR (extrapolated at U_{crit}) and aerobic scope (MMR - SMR). The resulting metabolic rate values were then normalized to a constant mass of 1 g by using the scaling constants and coefficients derived in Fig S4.




Figure S4. Determination of SMR and MMR mass scaling constants (y-intercept, a) and coefficients (slope, b) for juvenile mahi-mahi used to normalize metabolic rate measurements to a 1 g fish. Least-squares linear regressions of SMRs (open circles) and MMRs (filled circles) obtained by swimming respirometry were used to derive the following values, respectively: $b = 0.68$, $\log a = -2.03$; $b = 0.86$, $\log a = -2.26$. Adjusted SMR and MMR values were calculated for each individual by adding the difference of the corresponding residual from each regression line to the predicted SMR or MMR value for a 1 g fish.




Figure S5. Raw data (not scaled for mass) comparisons of (A) standard metabolic rate (SMR), (B) maximum metabolic rate (MMR), (C) aerobic scope and (D) cost of transport at U_{crit} from swimming respirometry experiments of juvenile mahi-mahi exposed to 4.2, 17 or 30 $\mu\text{g L}^{-1}$ ΣPAHs ($n = 7, 9$ and 8 , respectively) for 24-h prior to swimming (shaded) or to 1.2 $\mu\text{g L}^{-1}$ ΣPAHs ($n = 5$) for 48-h initiated <12 hpf (hatched bar) and then raised for ~25 d to the juvenile stage. The control group represents fish from all 4 control cohorts combined ($n = 21$). Data are presented as means \pm SEM. *Significantly different from controls by one-way ANOVA and Holm-Sidak multiple comparison procedure ($P < 0.05$).

